

# Τίτλος άρθρου: *Ανθεκτικότητα της πράσινης αφίδας της ροδακινιάς σε εντομοκτόνα την τελευταία εικοσαετία*

Συγγραφείς:

Ιωάννης Τ. Μαργαριτόπουλος<sup>1</sup>, Αμαλία Ν. Κατή<sup>3</sup>, Κωνσταντίνος Χ. Βουδούρης<sup>1</sup>, Παναγιώτης Ι. Σκούρας<sup>2</sup>, & Ιωάννης Α. Τσιτσιπής<sup>4</sup>

<sup>1</sup> Τμήμα Φυτοπροστασίας Βόλου, Ινστιτούτο Βιομηχανικών & Κτηνοτροφικών Φυτών, Ελληνικός Γεωργικός Οργανισμός «ΔΗΜΗΤΡΑ», Οδός Φυτόκου 47, 380 01 Βόλος. Επικοινωνία: Τηλ.: 2421061087, Fax: 2421061038, email: johnmargaritopoulos@gmail.com

<sup>2</sup> Εργαστήριο Εντομολογίας & Γεωργικής Ζωολογίας, Τμήμα Τεχνολόγων Γεωπόνων, Σχολή Τεχνολογίας Γεωπονίας & Τεχνολογίας Τροφίμων και Διατροφής, ΤΕΙ Πελοποννήσου, 241 00 Αντικάλamos, Μεσσηνία

<sup>3</sup> Εργαστήριο Εφαρμοσμένης στη Γεωργία Ζωολογίας και Παρασιτολογίας, Τμήμα Γεωπονίας, Σχολή Γεωπονίας, Δασολογίας και Φυσικού Περιβάλλοντος, Αριστοτέλειο Πανεπιστήμιο Θεσσαλονίκης, 541 24 Θεσσαλονίκη

<sup>4</sup> Εργαστήριο Εντομολογίας, Τμήμα Γεωπονίας Φυτικής Παραγωγής και Αγροτικού Περιβάλλοντος, Πανεπιστήμιο Θεσσαλίας, Οδός Φυτόκου, 384 46 Βόλος. Παρούσα Διεύθυνση: Μαινάλου 4, 152 35 Βρυλίσσια

## ΒΙΒΛΙΟΓΡΑΦΙΑ

1. Bass C, Puinean AM, Zimmer CT, Denholm I, Field LM, Foster SP, et al., The evolution of insecticide resistance in the peach potato aphid, *Myzus persicae*, *Insect Biochem Mol Biol* 51:41–51 (2014).
2. Bass C, Denholm I, Williamson MS, and Nauen R, The global status of insect resistance to neonicotinoid insecticides, *Pestic Biochem Physiol* 121:78–87 (2015).
3. Sparks TC and Nauen R, IRAC: Mode of action classification and insecticide resistance management, *Pestic Biochem Physiol* 121:122–128 (2015).
4. Cox D, Denholm I, and Devonshire A, Monitoring of insecticide resistance in *Myzus persicae* from Greece, ed. by Simon, J.-C., Dedryver, C.A., Rispe, C., and Hullé, M., *Aphids in a new millennium*, INRA Editions, Paris, pp. 275–280 (2004).
5. Anstead JA, Mallet J, and Denholm I, Temporal and spatial incidence of alleles conferring knockdown resistance to pyrethroids in the peach-potato aphid, *Myzus persicae* (Hemiptera: Aphididae), and their association with other insecticide resistance mechanisms, *Bull Entomol Res* 97:243–252 (2007).
6. Margaritopoulos JT, Skouras PJ, Nikolaidou P, Manolikaki J, Maritsa K, Tsamandani K, et al., Insecticide resistance status of *Myzus persicae* (Hemiptera: Aphididae) populations from peach and tobacco in mainland Greece, *Pest Manag Sci* 63:821–829 (2007).
7. Margaritopoulos JT, Tsamandani K, Kanavaki OM, Katis NI, and Tsi-tsipis JA, Efficacy of pymetrozine against *Myzus persicae* and in reducing potato virus Y transmission on tobacco plants, *J Appl Entomol* 134:323–332 (2010).
8. Kati AN, Mandrioli M, Skouras PJ, Malloch GL, Voudouris CC, Ventur-elli M, et al., Recent changes in the distribution of carboxylesterase genes and associated chromosomal rearrangements in Greek populations of the tobacco aphid *Myzus persicae nicotianae*, *Biol J Linn Soc* 113:455–470 (2014).
9. Voudouris CC, Kati AN, Sadikoglou E, Williamson M, Skouras PJ, Dimotsiou O, et al., Insecticide resistance status of *Myzus persicae* in Greece: long-term surveys and new diagnostics for resistance mechanisms, *Pest Manag Sci* 72:671–683 (2016).
10. Voudouris CC, Williamson MS, Skouras PJ, Kati AN, Sahinoglou AJ, and Margaritopoulos JT, Evolution of imidacloprid resistance in *Myzus persicae* in Greece and susceptibility data for spirotetramat, *Pest Manag Sci* 73:1804–1812 (2017).
11. Cutler P, Slater R, Edmunds AJF, Maienfisch P, Hall RG, Earley FGP, et al., Investigating the mode of action of sulfoxaflor: a fourth-generation neonicotinoid, *Pest Manag Sci* 69:607–619 (2013).
12. Scott JG and Wen Z, Cytochromes P450 of insects: the tip of the iceberg, *Pest Manag Sci* 57:958–967 (2001).

# Τίτλος άρθρου: Χειμερινός λήθαργος φυλλοβόλων οπωροφόρων και ακρόδρυων δένδρων

## 2. Μοντέλα υπολογισμού και δεδομένα συσσώρευσης ψύχους σε περιοχές της Ελλάδας και Κύπρου

Συγγραφείς: Δρ. Δρογούδη Παυλίνα<sup>1</sup>, Δρ. Παντελίδης Γεώργιος<sup>1</sup> και Ομ. καθ. Βασιλακάκης Μιλτιάδης<sup>2</sup>

<sup>1</sup> Τμήμα Φυλλοβόλων Οπωροφόρων Δένδρων, Ινστιτούτο Γενετικής Βελτίωσης & Φυτογενετικών Πόρων, ΕΛΓΟ 'ΔΗΜΗΤΡΑ', Σ.Σ. Νάουσας 38, Νάουσα, [www.pomologyinstitute.gr](http://www.pomologyinstitute.gr)

<sup>2</sup> Αριστοτέλειο Πανεπιστήμιο Θεσσαλονίκης, Τμήμα Γεωπονίας, Εργαστήριο Δενδροκομίας

### ΒΙΒΛΙΟΓΡΑΦΙΑ

- Aslani Aslamarz A.A., K. Vahdati, M. Rahemi, D. Hassani 2009. Estimation of the chilling and heat requirement of some Persian walnut cultivars. *HortScience*, 44: 697-701.
- Alburquerque, N., F. Garcia-Montiel, et al., 2008. Chilling and heat requirements of sweet cherry cultivars and the relationship between altitude and the probability of satisfying the chill requirements. *Envir. Exper. Bot.* 64: 162-170.
- Atkinson, C.J., Brennan, R.M., Jones, H.G., 2013. Declining chilling and its impact on temperate perennial crops. *Envir. Exper. Bot.* 91: 48-62.
- Baldocchi, D., Wong, S., 2008. Accumulated winter chill is decreasing in the fruit growing regions of California. *Clim. Change* 87: S153-S166.
- Blanke, M.M., and Kunz, A. 2017. Cherry phenology as a bioindicator for climate change. *Acta Hort.* 1162: 1-8.
- Campoy et al., 2012. The fulfillment of chilling requirements and the adaptation of apricot (*Prunus armeniaca* L.) in warm winter climates: An approach in Murcia (Spain) and the Western Cape (South Africa). *Eur. J. Agronomy* 37: 43-45.
- Charrier G, Bonhomme M, Lacoite A, Améglio T. 2011. Are budburst dates, dormancy and cold acclimation in walnut trees (*Juglans regia* L.) under mainly genotypic or environmental control? *Int J Biometeorol.* 55:763-74.
- Costa, C., Stassen, P.J.C., Mudzunga, J., 2004. Chemical rest breaking agents for the South African pome and stone fruit industry. *Acta Hort.* 636: 295-302.
- Darbyshire, R., Webb, L., Goodwin, I., Barlow, S., 2011. Winter chilling trends for deciduous fruit trees in Australia. *Agricultural and Forest Meteorology.* 151(8): 1074-1085.
- Dennis, F.G., 2003. Problems in standardizing methods for evaluating the chilling requirements for the breaking of dormancy in buds of woody plants. *HortSci.* 38: 347-350.
- Drogoudi P., K. Kazantzis & M.M. Blanke 2017. Climate change effects on cherry flowering in Northern Greece. *Acta Hort.* 1162: 45-49.
- Egea, J., Ortega, E., Martínez-Gómez, P., Dicenta, F., 2003. Chilling and heat requirements of almond cultivars for flowering. *Environmental and Experimental Botany.* 50(1): 79-85.
- Elloumi, O., Ghrab, M., Kessentini, H., Ben Mimoun, M., 2013. Chilling accumulation effects on performance of pistachio trees cv. Ma-teur in dry and warm area climate. *Sci. Hort.* 159: 80-87.
- Erez, A. 2000. Bud dormancy; phenomenon, problems and solutions in the tropics and subtropics. Temperate fruit crops in warm climates. A. Erez. Dordrecht, The Netherlands, Kluwer Academic Publishers: 17-48.
- Erez et al., 1990. The dynamic model for rest completion in peach buds. *Acta Hort.* 276: 165-174.
- Erez, A., 1995. Means to compensate for insufficient chilling to improve bloom and leafing. *Acta Hort.* 395: 81-95.
- Fishman, S., Erez, A., Couvillon, G.A., 1987. The temperature dependence of dormancy breaking in plants—mathematical analysis of a two-step model involving a cooperative transition. *J. Theor. Biol.* 124: 473-483.
- Funes et al., 2016. Future climate change impacts on apple flowering date in a Mediterranean subbasin. *Agr. Water Manag.* 164: 19-27.
- Garbone and Schwartz, 1993. Potential impact of winter temperature increases on South Carolina peach production. *Climat Res.* 2: 225-233.
- Ghrab, M., Ben Mimoun, M., Masmoudi, M.M., Ben Mechlia, N., 2014. The behaviour of peach cultivars under warm climatic conditions in the Mediterranean area. *Int. J. Env. St.*, 71: 3-14.
- Kolářová, E., Nekovář, J., Adamík, P., 2014. Long-term temporal changes in central European tree phenology (1946-2010) confirm the recent extension of growing seasons. *International Journal of Biometeorology.* 58(8): 1739-1748.
- Lammerts, W.E., 1941. An evaluation of peach and nectarine varieties in terms of winter chilling requirements and breeding possibilities. *P. Am. Soc. Hortic. Sci.* 39: 205-211.
- Linsley-Noakes, G. C. and P. Allan 1994. Comparison of two models for the prediction of rest completion in peaches. *Sci. Hort.* 59: 107-113.
- Luedeling et al. 2009. Validation of winter chill models using historic records of walnut phenology. *Agric. For Meteorol.* 149: 1854-1864.
- Luedeling, E., Brown, P.H., 2011. A global analysis of the comparability of winter chill models for fruit and nut trees. *Int. J. Biometeorol.* 55: 411-421.
- Luedeling, E., L. Guo, et al., 2013. "Differential responses of trees to temperature variation during the chilling and forcing phases." *Agric. Forest Met.* 181: 33-42.
- Luedeling, E., M. Zhang, et al., 2009. Validation of winter chill models using historic records of walnut phenology. *Agric. Forest Met.* 149: 1854-1864.
- Luedeling, E., Zhang, M., Girvetz, E.H., 2009a. Climatic changes lead to declining winter chill for fruit and nut trees in California during 1950-2009. *PLoS One*, 4, e6166.
- Luedeling, E., M. Zhang, G. McGranahan, C. Leslie, 2009b. Validation of winter chill models using historic records of walnut phenology. *Agric Forest Meteorology* 149: 1854-1864.
- Luedeling, E., Blanke, M., Gebauer, J., 2015. Chilling Challenges in a Warming World. *Acta Hort.* 1099: 901-908.
- Miranda (2013) Evaluation and fitting of models for determining peach phenological stages at a regional scale. *Agric Forest Meteo* 178-179: 129-139.
- Okie, 1998. Handbook of peach and nectarine varieties. *Agr. Handbook No. 714*, USA.
- Palasciano M, and Gaeta L. 2017. Comparison of different models for chilling requirements evaluation of sweet cherry cultivars in a Mediterranean area. *Acta Hort.* 1161: 405-410.
- Palasciano, M., Gaeta, L., 2017. Comparison of different models for chilling requirements evaluation of sweet cherry cultivars in a Mediterranean area. *Acta Hort.* 1161: 405-410.
- Ramirez, L., K. X. Sagredo, et al. 2010. Prediction models for chilling and heat requirements to estimate full bloom of almond cultivars in the Central Valley of Chile. *Acta Hort.* 872: 107-112.
- Richardson, E.A., Seeley, S.D., Walker, D.R., 1974. A model for estimating the completion of rest for Redhaven and Elberta peach trees. *HortSci.* 9: 331-332.
- Ruiz, D., J. A. Campoy, et al. 2007. Chilling and heat requirements of apricot cultivars for flowering. *Envir. Exper. Botany* 61: 254-263.
- Segura et al., 2017. Late-blooming in almond: A controversial objective. *Sci. Hort.* 224: 61-67.

39. UC Davis. Fruit and Nut research and information center. Dynamic Model & Chill Accumulation [http://ucanr.edu/sites/fruittree/how-to\\_guides/dynamic\\_model\\_-\\_chill\\_accumulation/](http://ucanr.edu/sites/fruittree/how-to_guides/dynamic_model_-_chill_accumulation/)
40. Viti, R., L. Andreini, et al., 2010. Effect of climatic conditions on the overcoming of dormancy in apricot flower buds in two Mediterranean areas: Murcia (Spain) and Tuscany (Italy). *Sci. Hortic.* 124: 217-224.
41. Wall, C., Dozier, W., Ebel, R.C., Wilkins, B., Woods, F., Foshee III, W., 2008. Vegetative and floral chilling requirements of four new kiwi cultivars of *Actinidia chinensis* and *A. deliciosa*. *HortSci.* 43: 644-647.
42. Weinberger, J.H., 1950. Chilling requirements of peach varieties. *P. Am. Soc. Hortic. Sci.* 56: 122-128.
43. Vahdati K, A. Aslani Aslamarz, M. Rahemi, D. Hassani, C. Leslie, 2012. Mechanism of seed dormancy and its relationship to bud dormancy in Persian walnut. *Environmental and Experimental Botany* 75: 74–82.
44. USDA Risk Management Agency Valdosta Regional Office, 2017. Peach Variety Listing Table, 2017. [https://www.rma.usda.gov/fields/ga\\_rso/2017/peachvl.html](https://www.rma.usda.gov/fields/ga_rso/2017/peachvl.html)

# Τίτλος άρθρου: Η βρωμούσα *Halyomorpha halys* στην Ελλάδα

Συγγραφείς: Εμμανουήλ Ναβροζίδης<sup>1</sup>, Στέφανος Ανδρεάδης<sup>2</sup>, Αργύρης Φαρμάκης<sup>1</sup> και Αθανάσιος Πισαλίδης<sup>3</sup>

<sup>1</sup> Αλεξάνδρειο Τεχνολογικό Εκπαιδευτικό Ινστιτούτο, Σχολή Τεχνολογίας Γεωπονίας και Τεχνολογίας Τροφίμων και Διατροφής, Τμήμα Τεχνολόγων Γεωπόνων, 57400 Σίνδος

<sup>2</sup> ΕΛ.Γ.Ο. – “ΔΗΜΗΤΡΑ”, Ινστιτούτο Γενετικής Βελτίωσης και Φυτογενετικών Πόρων 57001 Θέρμη

<sup>3</sup> Νέα Έφρεσος, 60100 Πιερία

## ΒΙΒΛΙΟΓΡΑΦΙΑ

1. Hoebeke, E.R., Carter, M.E. 2003. *Halyomorpha halys* (Stål) (Heteroptera: Pentatomidae): A polyphagous plant pest from Asia newly detected in North America. *Proceedings of the Entomological Society of Washington* 105: 225–237.
2. Xu, J., Fonseca, D.M., Hamilton, G.C., Hoelmer, K.A., Nielsen, A.L. 2014. Tracing the origin of US brown marmorated stink bugs, *Halyomorpha halys*. *Biological Invasions* 16: 153–166.
3. Haye, T., Abdallah, S., Garipey, T., Wyniger, D. 2014. Phenology, life table analysis, and temperature requirements of the invasive brown marmorated stink bug, *Halyomorpha halys*, in Europe. *Journal of Pest Science* 87: 407–418.
4. Wermelinger, B., Wyniger, D., Forster, B. 2008. First records of an invasive bug in Europe *Halyomorpha halys* Stål (Heteroptera: Pentatomidae), a new pest on woody ornamentals and fruit trees? *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 81: 1–8.
5. Bergmann, E., Bernhard, K.M., Bernon, G., Bickerton, M., Gill, S., Gonzales, C., Hamilton, G.C., Hedstrom, C., Kamminga, K., Koplinka-Loehr, C., Krawczyk, G., Kuhar, T.P., Kunkel, B., Lee, J., Leskey, T.C., Martinson, H., Nielsen, A.L., Raupp, M., Shearer, P., Shrewsbury, P., Walgenbach, J., Whalen, J., Wiman N. 2013: *Host plants of the brown marmorated stink bug in the U.S. National Institute of Food and Agriculture U.S. Department of Agriculture, Specialty Crop Research Initiative under award numbers 2016-51181-25409 and 2011-51181-30937.* ([http://www.stopbmsb.org/where-is-bmsb/host-plants/#host\\_plants\\_table](http://www.stopbmsb.org/where-is-bmsb/host-plants/#host_plants_table))
6. Lee, D.H., Wright, S.E., Boiteau, G., Vincent, C., Leskey, T.C. 2013. Effectiveness of glues for harmonic radar tag attachment on *Halyomorpha halys* (Hemiptera: Pentatomidae) and their impact on adult survivorship and mobility. *Environmental Entomology* 42: 515–523.
7. Lee, D.H. 2015. Current status of research progress on the biology and management of *Halyomorpha halys* (Hemiptera: Pentatomidae) as an invasive species. *Applied Entomology and Zoology* 50: 277–290.
8. Rice, K.B., Bergh, C.J., Bergmann, E.J., Biddinger, D.J., Dieckhoff, C., Dively, G., Fraser, H., Garipey, T., Hamilton, G., Haye, T., Herbert, A., Hoelmer, K., Hooks, C.R., Jones, A., Krawczyk, G., Kuhar, T., Martinson, H., Mitchell, W., Nielsen, A.L., Pfeiffer, D.G., Raupp, M.J., Rodriguez-Saona, C., Shearer, P., Shrewsbury, P., Venugopal, D.P., Whalen, J., Wiman, N.G., Leskey, T.C., Tooker, J.F. 2014. Biology, ecology, and management of brown marmorated stink bug (Hemiptera: Pentatomidae). *Journal of Integrated Pest Management* 5: 1–13.
9. Bariselli, M., Bugiani, R., Maistrello, L. 2016. Distribution and damage caused by *Halyomorpha halys* in Italy. *OEPP/EPPO Bulletin* 46: 332–334.
10. Lupi, D., Dioli, P., Limenta, L. 2017. First evidence of *Halyomorpha halys* Stål (Hemiptera, Heteroptera, Pentatomidae) feeding on rice (*Oryza sativa* L.). *Journal of Entomological and Acarological Research* 49: 67–71.
11. Bernardinelli, I., Malossini, G., Benvenuto, L. 2017. *Halyomorpha halys*: risultati preliminari di alcune attività sperimentali condotte in Friuli Venezia Giulia nel 2016. *Notiziario ERSA* 1: 24–26 [in Italian].
12. Pasqualini, E., Burgio, G., Preti, M., Depalo, L., Masetti, A. 2017. Emergenza cimice asiatica: approccio integrato per la difesa. *Frutticoltura* 3: 24-27 [in Italian].
13. Milonas, P.G., Partsinevelos, G.K. 2014. First report of brown marmorated stink bug *Halyomorpha halys* Stål (Hemiptera: Pentatomidae) in Greece. *OEPP/EPPO Bulletin* 44: 183–186.
14. Garipey, T.D., Bruin, A., Haye, T., Milonas, P., Véték, G. 2015. Occurrence and genetic diversity of new populations of *Halyomorpha halys* in Europe. *Journal of Pest Science* 88: 451–460.
15. Morrison III, W.R., Milonas, P., Kapantaidaki, D.E., Cesari, M., Di Bella, E., Guidetti, R., Haye, T., Maistrello, L., Moraglio, S.T., Piemontese, L., Pozzebon, A., Ruocco, G., Short, B.D., Tavella, L., Véték, G., Leskey, T.C. 2017. Attraction of *Halyomorpha halys* (Hemiptera: Pentatomidae) haplotypes in North America and Europe to baited traps. *Scientific Reports* 7: 16941.
16. Μυλωνάς, Π.Γ., Παρτσινέβελος, Γ.Κ., Καπανταϊδάκη Δ.Ε., Μιχαηλάκης Α. 2017. Το έντομο *Halyomorpha halys*: Ένας νέος σημαντικός εχθρός για τις καλλιέργειες στην Ελλάδα. *Γεωργία-Κτηνοτροφία* 9: 54–56.
17. Andreadis, S.S., Navrozidis, E.I., Farmakis, A., Pisalidis, A. 2018. First Evidence of *Halyomorpha halys* (Hemiptera: Pentatomidae) Infesting Kiwi Fruit (*Actinidia chinensis*) in Greece. *Journal of Entomological Science* (in press).
18. [FAO] Food and Agriculture Organization of the United Nations. 2018. FAOSTAT, Statistics Division: Crops. URL: <http://www.fao.org>.