

# Ο ρόλος των παρεμποδιστών νιτροποίησης στην σύγχρονη γεωργία

Ευαγγελία Παπαδοπούλου, Δημήτρης Καρούζας

Πανεπιστήμιο Θεσσαλίας, Τμήμα Βιοχημείας και Βιοτεχνολογίας,  
Εργαστήριο Βιοτεχνολογίας Φυτών και Περιβάλλοντος,  
Βιόπολις, 41500, Λάρισα. Email [evaparadopoulos@bio.uth.gr](mailto:evaparadopoulos@bio.uth.gr);  
[dkarouzas@bio.uth.gr](mailto:dkarouzas@bio.uth.gr)

## ΒΙΒΛΙΟΓΡΑΦΙΑ

1. O'Brien et al. 2016 *Molecular Plant*. doi:10.1016/j.molp.2016.05.004
2. Raun and Johnson 1999 *Agronomy J* 9: 357-363
3. Coskun et al. 2017 *Nature Plants* 3:17074
4. van Groenigen et al. 2015 *Soil* 1:235-256
5. Abalos et al. 2019 *J Appl Ecol*56:2454-2466
6. Beeckman et al. 2018 *Curr Opin Biotechnol* 50: 166-173
7. Venter et al. 2004 *Science* 304:66-74
8. Leininger et al. 2006 *Nature* 442:806-809
9. Prosser and Nicol 2008 *Environ Microbiol*10:2931-2941
10. Gubry-Rangin et al. 2010 *FEMS Microbiol Ecol* 74: 566-574 11. Zhang et al. 2010 *PNAS USA* 107: 17240-17245
11. Tourna et al. 2011 *Proc Natl Acad Sci U S A* 108: 8420-8425
12. Daims et al. 2015 *Nature* 528: 504-509
13. van Kessel et al. 2015 *Nature* 528:555-559
14. Li et al. 2019 *Soil Biol Biochem* 138:107609
15. Wang et al. 2019 *Soil Biol Biochem* 135:392-395
16. Feng et al. 2016 *Agriculture Ecosystem and Environment* 231:218-228
17. 18. Subbarao et al. 2012 *Advances in Agronomy*114:250-272
18. Mc Carty 1999 *Biology and Fertility of Soils* 29:1-9
19. Solansky 1982 *Blickfeld* 61: 1-4
20. Goring 1962 *Soil Sci* 93: 211-218
21. Zerulla et al. 2001 *Biol Fertil Soils* 34:79-84
22. Di and Cameron 2011 *Journal of Soils Sediments* 11:1032-1039
23. Kleineidam et al. 2011 *Chemosphere* 84:182-186
24. Cui et al. 2013 *Soil Biol Biochem* 66:214-221
25. Baszczyk et al. 2013 *Int J Food Science*<http://dx.doi.org/10.1155/2013/585931>
26. Papadopoulou et al. 2016 *Appl Environ Microbiol*82:747-755
27. Papadopoulou et al. 2020 <https://doi.org/10.1101/2020.04.07.023168>
28. Fabian et al. 2014 *Applied Geochemistry* 48: 207-216
29. Philippot et al. 2011 *Global Change Biology*17:1497-1504

# Επιδράσεις της κλιματικής αλλαγής στην παραγωγή κερασιών και στρατηγικές προσαρμογής

Π. Δρογούδη<sup>1</sup>, Κ. Καζαντζής<sup>1</sup>, Γ. Παντελίδης<sup>1</sup>, A. Kunz<sup>2</sup>, M. Blanke<sup>2</sup>

<sup>1</sup>Τμήμα Φυλλοβόλων Οπωροφόρων Δένδρων (ΤΦΟΔ) Νάουσας, Ινστιτούτο Γενετικής Βελτίωσης και Φυτογενετικών Πόρων,

Ελληνικός Γεωργικός Οργανισμός 'Δήμητρα', Σ.Σ. Νάουσας 38, Τ.Κ. 59035 Νάουσα

<sup>2</sup>INRES-Horticultural Science, University of Bonn, Auf dem Hóegel 6, 53121 Bonn, Germany

## ΒΙΒΛΙΟΓΡΑΦΙΑ

1. Blanke M.M., Kunz A. (2009a). Effects of climate change on pome fruit phenology at Klein-Altendorf [Auswirkungen rezenter Klimaveränderungen auf die Phanologie von Kernobst am Standort Klein-Altendorf]. *Erwerbs-Obstbau* 51:101–114 (special edition on climate change).
2. Blanke M.M., Kunz A. (2009b). Misconceptions about the effects of climate change on horticulture. *Proc Benelux Horticult Soc Annu Meeting, Gembloux, Belgium*, p 12. <http://www.beneluxshs.eu/index.html>. Accessed 3 Apr 2009.
3. Kaufmann H., Blanke M.M. (2017). Performance of three numerical models to assess winter chill in fruit trees—a case study using cherry as model crop in Germany. *Reg Environ Change* 17:1–9.
4. Kaufmann H., Blanke M.M. (2018). Substitution of winter chilling by spring forcing for flowering of sweet cherry? *Sci Hortic (Amsterdam)* 244:75–81.
5. Kaufmann H., Blanke M.M. (2019). Chilling requirements of Mediterranean fruit crops in a changing climate. *Acta Hort* 1242:275–280.
6. Luedeling E., Kunz A., Blanke M.M. (2011). More chilling for fruit trees in warmer winters [Mehr Chilling für Obstbaume in warmeren Wintern]? *Erwerbs-Obstbau* 53:145–155.
7. Luedeling E., Kunz A., Blanke M.M. (2013). Identification of chilling and heat requirements of cherry trees—a statistical approach. *Int J Biometeorol* 57:679–689.
8. Wenden B. et al. (2017). Harmonisation of phenology stages. *Acta Hort* 1162:9–12.
9. Xoplaki E. (2020). Climate and environmental changes in the Med region (MEDECC). *Euro-Mediterr J Environ Integr* (volume: 5).
10. Χατζηχαρίσης Ι., Καζαντζής Κ. (2014). Η Κερασιά και η καλλιέργειά της. Εκδόσεις ΑγροΤύπος, σελίδες 440.

# Βιολογική καταπολέμηση της Μύγας της Μεσογείου με Εντομοπαθογόνους νηματώδεις

Απόστολος Καπράνας, Άννα Χρονοπούλου,  
Ιωάννα Λύτρα, Δημήτρης Παπαχρήστος  
και Παναγιώτης Μυλωνάς

Τμήμα Εντομολογίας και Γεωργικής Ζωολογίας,  
Μπενάκειο Φυτ. ΙνστιτούτοΣτ. Δέλτα 8 Κηφισιά  
a.kapranas@bpi.gr

## ΒΙΒΛΙΟΓΡΑΦΙΑ

1. Campbell, J. F., Lewis, E. E., Stock, S. P., Nadler, S. and Kaya, H.K. (2003). Evolution of host search strategies in entomopathogenic nematodes. *Journal of Nematology* 35: 142–145
2. Campbell, J. and Gaugler, R. (1993). Nictation behaviour and its ecological implications in the host search strategies of entomopathogenic nematodes (Heterorhabditidae and Steinernematidae). *Behaviour* 126: 154–169.
3. Campbell, J. F. and Kaya, H. K. (1999). How and why a parasitic nematode jumps. *Nature* 397: 485–486.
4. Campos-Herrera R (2015) *Nematode Pathogenesis of Insects and other Pests - Ecology and Applied Technologies for Sustainable Plant and Crop protection*. Springer International Publishing
5. Dembilio, O., Llacer, E., De Altube, M. M. and Jacas, J. A. (2010). Field efficacy of imidacloprid and *Steinernema carpocapsae* in a chitosan formulation against the red palm weevil *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae) in Phoenix canariensis. *Pest Management Science* 66: 365–370.
6. Dillman, A.R. and Sternberg, P.W. (2012). Quick guide: entomopathogenic nematodes. *Current Biology* 22: 430–431.
7. Gazit, Y., Rossler, Y., and Glazer, I. (2000). Evaluation of entomopathogenic nematodes for the Control of Mediterranean Fruit Fly (Diptera: Tephritidae). *Biocontrol Science and Technology* 10: 157–164.
8. Garrido-Jurado, I., Torrent, J., Barrón, V., Corpas, A., Quesada-Moraga, E. (2011). Soil properties affect the availability, movement, and virulence of entomopathogenic fungi conidia against puparia of *Ceratitis capitata* (Diptera: Tephritidae). *Biological Control* 58: 277–285.
9. Garriga, A., Morton, A., Ribes, A., Garcia-del-Pino, F. (2020). Soil emergence of *Drosophila suzukii* adults: A susceptible period for entomopathogenic nematodes infection. *J. Pest Science* 93: 639–646.
10. Grewal, P. S., Lewis, E. E., Gaugler, R. and Campbell, J. F. (1994). Host finding behaviour as a predictor of foraging strategy in entomopathogenic nematodes. *Parasitology* 108, 207–215.
11. Griffin, C. (2015). Behaviour and population dynamics of entomopathogenic nematodes following application. *Nematode Pathogenesis of Insects and Other Pests – Ecology and Applied Technologies for Sustainable Plant and Crop Protection*. Springer International Publishing, pp. 57–96.
12. Grewal, P.S., Selvan, S., & Gaugler, R. (1994). Thermal adaption of entomopathogenic nematodes: niche breadth for infection, establishment, and reproduction. *Journal of Thermal Biology* 19: 245–253.
13. Kapranas, A., Malone, B., Quinn, S., McNamara, L., Williams, C. D., O'Tuama, P., Peters, A. and Griffin, C. T. (2017) Efficacy of entomopathogenic nematodes for control of large pine weevil, *Hyllobius abietis*: effects of soil type, pest density and spatial distribution. *J Pest Science* 90: 495–505
14. Kapranas, A., Maher, A.M.D., and Griffin, C. T. (2017) The influence of organic matter content and media compaction on the dispersal of entomopathogenic nematodes with different foraging strategies. *Parasitology* 144: 1956–1963.
15. Kaya, H.K., and Gaugler, R. (1993). Entomopathogenic nematodes. *Annual Review Entomology* 38: 181–206.
16. Lacey, L. A. and Georgis, R. (2012) Entomopathogenic nematodes for control of insect pests above and below ground with comments on commercial production. *J Nematology* 44: 218–225.
17. Lewis, E. E., Campbell, J., Griffin, C., Kaya, H. and Peters, A. (2006). Behavioral ecology of entomopathogenic nematodes. *Biological Control* 38: 66–79.
18. Martinez de Altube, M. D. M., Strauch, O., De Castro, G. F. and Pena, A. M. (2008). Control of the flat headed root borer *Capnodis tenebrionis* (Linné) (Coleoptera: Buprestidae) with the entomopathogenic nematode *Steinernema carpocapsae* (Weiser) (Nematoda: Steinernematidae) in a chitosan formulation in apricot orchards. *BioControl* 53: 531–539.
19. Shapiro-Ilan, D. I., Han, R. and Dolinski, C. (2012). Entomopathogenic nematode production and application technology. *J. Nematology* 44: 226–235.
20. Shapiro-Ilan, D. I., Gouge, D. H., Piggott, S. J., Fife, J. P. (2006). Application technology and environmental considerations for use of entomopathogenic nematodes in biological control. *Biological Control* 38: 124–133
21. Stark, J. D. and Vargas, R (2009). An evaluation of alternative insecticides to diazinon for control of Tephritid fruit flies (Diptera: Tephritidae) in soil. *J Economic Entomology* 102: 139–143.
22. Τζανακάκης, Μ.Ε., και Κατσόγιαννος, Β.Ι. (1998). Έντομα Καρποφόρων Δέντρων Και Αμπέλου. Αθήνα, Εκδόσεις ΑγροΤύπος, 359 σελ.