

## ΔΕΝΔΡΟΚΟΜΙΑ

### Επίδραση του βιοδιεγέρτη Bud14 στην καλλιέργεια της ακτινιδιάς

#### Πώς επηρεάζει το βάρος των καρπών και τη συνολική παραγωγή

##### Βιβλιογραφία

- Costa G, Vizzotto G and O. Lain. 1995. Fruiting performance of kiwifruit cv. Hayward affected by use of hydrogen cyanamide *Acta Hort.* 444: 473-478.
- Costa, G., G. Vizzotto, and O. Lain. 1997. Fruiting performance of kiwifruit cv. Hayward affected by use of hydrogen cyanamide. *Acta Hort.* 444:473-478.
- Cruz-Castillo, J., Woolley, D., anf G. Lawes. (2002). Kiwifruit size and CPPU response are influenced by the time of anthesis. *Sci. Hort.* 95: 23–30.
- Engin, H, Gökbayrak, Z. and A. Dardeniz (2010). Effects of Hydrogen Cyanamide on the Floral Morphogenesis of Kiwifruit Buds. *Chilean J. Agric. Res.* 70. 503-509.
- Guedon Y, Bhartlemy D, Caraglio Y and E. Costes (2001). Pattern analysis in branching and axillary flowering sequences. *Journal Theor. Biol.* 212: 481-520.
- Lai, R., Woolley, D. and G. Lowes. (1990). The effect of inter-fruit competition, type of fruiting lateral and time of anthesis on fruit growth of kiwifruit. *Aust. J. Agr. Res.* 65:87-96.
- Marques, L., Mello-Farias, P., Fernandes, R., Martins, C., Konzen L., and M. Malgarim. (2021). Alternative budburst inducers for kiwifruit vines grown in an organic system. *Rev. Bra. Frutic.* 43: 1-10.
- Mcpherson HG, Hall AJ and C. Stanley (1994). Seasonal and regional variation in bud break and flowering of kiwifruit vines in New Zealand. *New Zealand J. Crop Hortic. Sci.* 22: 263-276.
- Mcpherson, H., Richardson, A., Snelgar, W., Patterson, K, and M. Currie. (2001) Flower quality and fruit size in kiwifruit. *New Zealand J. Crop and Hortic. Sci.* 29:2, 93-101.
- Salinero, M.C., Vela, P., and

M. Sainz. (2009). Phenological growth stages of kiwifruit (*Actinidia deliciosa*'Hayward'). *Sci. Hort.* 121::27-31.

- Pang X, Halaly T, Crane O, Keilin T, Keren-Keiserman A, Ogrodotvitch A, Galbraith D, Or E. (2007). Involvement of calcium signalling in dormancy release of grape buds. *J Exp Bot.* 58:3249-62.
- Patterson, K., Snelgar, W., Richardson, A. and H. McPherson (1999). Flower quality and fruit size of Hayward kiwifruit. *Acta Hort.* 498: 143-150.
- Pichakum, A.; Chaiwimol, W.; Meetam, M.; and W. Songnuan. (2018). Responses of green kiwifruit grown in low-chill area to hydrogen cyanamide application. *Acta Hortic.*, 1206:97-104.
- Di Tommaso, G., Piaggesi, A. and Mills, R. (2011). Bud break: Safe and efficient method of overcoming winter dormancy – New Zealand experiences with 'HORT16A' and 'HAYWARD' kiwifruit. *Acta Hortic.* 913, 275-282.
- Wall, C., Dozier, W., Ebel, R.C., Wilkins, B., Woods, F., and W. Foshee III. (2008). Vegetative and floral chilling requirements of four new kiwi cultivars of *Actinidia chinensis* and *A. deliciosa*. *HortSci.* 43: 644-647.

## ΦΥΤΑ ΜΕΓΑΛΗΣ

### ΚΑΛΛΙΕΡΓΕΙΑΣ

#### Κόκκινο ρύζι (*Oryza sativa* L.)

##### Γιατί είναι το πιο επιζήμιο και δύσκολο στην αντιμετώπιση ζιζάνιο στο ρύζι

##### Βιβλιογραφία

- Agnoun, Y., S.S.H. Biaou, M. Siè, R.S. Vodouhé and A. Ahanchédé. 2012. The African Rice *Oryza glaberrima* Steud: Knowledge Distribution and Prospects. *International Journal of Biology* 4(3):158-180.
- Burgos, N.R., V. Singh, T.M. Tseng, H. Black, N.D. Young, Z. Huang, K.E. Hyma, D.R. Gealy, and A.L. Caicedo. 2014. The impact of herbicide-resistant rice technology on
- phenotypic diversity and population structure of United States weedy rice. *Plant Physiology* 166:1208–1220.
- Chen, W., W. Wang, Y. Lyu, Y. Wu, P. Huang, S. Hu, X. Wei, G. Jiao, Z. Sheng, S. Tang, G. Shao, J. Luo. 2021. OsVP1 activates Sdr4 expression to control rice seed dormancy via the ABA signaling pathway. *The Crop Journal* 9:68-78.
- Cui, Y., B.K. Song, L.F. Li, Y.L. Li, Z. Huang, A.L. Caicedo, Y. Jia, and K.M. Olsen. 2016. Little white lies: pericarp color provides insights into the origins and evolution of Southeast Asian weedy rice. *G3: Genes, Genomes, Genetics* 6(12):4105-4114.
- Delouche, J.C., N.P. Burgos, R. Labrada, D.R. Gealy. 2007. Weedy rices: origin, biology, ecology and control. *Food and Agriculture Organization of the United Nations* 188:157 pages.
- Diarra, A., R.J. Smith, Jr., and R.E. Talbert. 1985. Growth and morphological characteristics of red rice (*Oryza sativa*) biotypes. *Weed Science* 33:310–314.
- Eleftherohorinos, I.G., K.V. Dhima and I.B. Vasilakoglou. 2002. Interference of red rice in rice grown in Greece. *Weed Science* 50:167-172.
- Engku, A.K., M. Norida, A.S. Juraimi, M.Y. Rafii, S.N.A. Abdullah, M.A. Alam. 2016. Gene flow from Clearfield® rice to weedy rice under field conditions. *Plant Soil Environment* 62:16–22.
- Fogliatto, S., A. Ferrero, F. Vidotto. 2020. How can weedy rice stand against abiotic stresses? A review. *Agronomy* 10:1284.
- Fogliatto, S., F. Vidotto, A. Ferrero. 2012. Morphological characterisation of Italian weedy rice (*Oryza sativa*) populations. *Weed Research* 52:60–69.
- Furukawa, T., M. Maekawa, T. Oki, I. Suda, S. Iida, H. Shimada, I. Takamure, and K. Kadokami. 2006. The Rc and Rd genes are involved in proanthocyanidin synthesis in rice pericarp. *The Plant Journal* 49:91-102.
- Gu, X.Y., M.E. Foley, D.P. Horvath, J.V. Anderson, J. Feng et al. 2011.



- Association between seed dormancy and pericarp color is controlled by a pleiotropic gene that regulates abscisic acid and flavonoid synthesis in weedy red rice. *Genetics* 189:1515–1524.
- Henry, H.J., N. Rice, D.L.E. Waters, S. Kasem, R. Ishikawa, Y. Hao, S. Dillon, D. Crayn, R. Wing, and D. Vaughan. 2010. Australian *Oryza*: Utility and Conservation. *Rice* 3:235–241.
  - Hua, L., D.R. Wang, L. Tan, Y. Fu, F. Liu, L. Xiao, Z. Zhu, Q. Fu, X. Sun, P. Gu, H. Cai, S.R. McCouch, and C. Suna. 2015. *LABA1*, a domestication gene associated with long, barbed awns in wild rice. *Plant Cell* 16, 2015; DOI: 10.1105/tpc.15.00260.
  - Huang, Z., S. Kelly, M. Matsuo, L-F Li, Y. Li, K.M. Olsen, Y. Jia, A.L. Caicedo. 2018. The role of standing variation in the evolution of weediness traits in South Asian weedy rice (*Oryza* spp.). *G3: Genes, Genomes, Genetics* 8 (11):3679-3690.
  - Huang, Z., N.D. Young, M. Reagon, K.E. Hyma, K.M. Olsen et al., 2017. All roads lead to weediness: Patterns of genomic divergence reveal extensive recurrent weedy rice origins from South Asian *Oryza*. *Molecular Ecology* 26:3151–3167.
  - Jabran, K. and M.N. Dogan. 2018. High carbon dioxide concentration and elevated temperature impact the growth of weeds but do not change the efficacy of glyphosate. *Pest Management Science* 74:766-771.
  - Jin, J., W. Huang, J-P. Gao, J. Yang, M. Shi, M-Z. Zhu, D. Luo, and H-X. Lin. 2008. Genetic control of rice plant architecture under domestication. *Nature Genetics* 40:1365–1369 doi:10.1038/ng.247.
  - Kaloumenos, N.S., Capote N., Aguado A., Eleftherohorinos I.G.. 2013. Red rice (*Oryza sativa*) cross-resistance to imidazolinone herbicides used in resistant rice cultivars grown in northern Greece. *Pesticide Biochemistry and Physiology* 105:177-183.
  - Kanapeckas, K.L., C.C. Vigueira, A. Ortiz, K.A. Gettler, N.R. Burgos A.J. Fischer, A.L. Lawton-Rauh. 2016. Escape to ferality: The endoferal origin of weedy rice from crop rice through dedomestication. *PLOS ONE*, DOI:10.1371/journal.
  - pone.0162676
  - Li, C., Zhou A., Sang, T. 2006. Rice domestication by reducing shattering. *Science* 311:1936–1939.
  - Liu, W., F. Ghouri, H. Yu, X. Li, S. Yu, M.Q. Shahid, X. Liu. 2017. Genome wide re-sequencing of newly developed Rice Lines from common wild rice (*Oryza rufipogon* Griff.) for the identification of NBS-LRR genes. *PLOS ONE* | <https://doi.org/10.1371/journal.pone.0180662>.
  - Londo, J.P. and B.A. Schaal. 2007. Origins and population genetics of weedy red rice in the USA. *Molecular Ecology* 16(21):4523-4535.
  - Matzrafi, M., C. Brunharo, P. Tehranchian, B.D. Hanson and M. Jasieniuk. 2019. Increased temperatures and elevated CO<sub>2</sub> levels reduce the sensitivity of *Conyza canadensis* and *Chenopodium album* to glyphosate. *Scientific Reports* 9:2228.
  - Mondal, T.K. and R.J. Henry. 2018. The wild *Oryza* genomes. *Compendium of Plant Genomes*. Springer International Publishing AG. 310p.
  - Noldin, J.A., J.M. Chandler, and G.N. McCauley. 2006. Seed longevity of red rice ecotypes buried in soil. *Planta Daninha* 24 (4): 611-620.
  - Pantone, D. J. and J. B. Baker. 1991. Weed–crop competition models and response. Surface analysis of red rice competition in cultivated rice: a review. *Crop Science* 31:1105–1110.
  - Prathepha, P. 2019. Complementary gene controlling the black hull phenotype and phenolics and flavonoids content in weedy rice (*Oryza sativa f. spontanea*) from northeastern Thailand. *Songklanakarin Journal of Science and Technology* 41:822-827.
  - Raza, A., A. Razzaq, S.S. Mahmood, X. Zou, X. Zhang, Y. Lv and J. Xu. 2019. Impact of climate change on crops adaptation and strategies to tackle its outcome: A review. *Plants* 8:34.
  - Reagon, M., C.S. Thurber, B.L. Gross, K.M. Olsen, Y. Jia, and A.L. Caicedo. 2010. Genomic patterns of nucleotide diversity in divergent populations of U.S. weedy rice. *BMC Evolutionary Biology* 10:180.
  - Shivrain V.K., N.R. Burgos, M.M.
  - Anders, S.N. Rajguru, J. Moore, M.A. Sales. 2007: Gene flow between Clearfield™ rice and red rice. *Crop Protection* 26: 349–356.
  - Shivrain V.K., N.R. Burgos, D.R. Gealy, K.A.K. Moldenhauer, C.J. Baquiereza. 2008. Maximum outcrossing rate and genetic compatibility between red rice (*Oryza sativa*) biotypes and Clearfield™ rice. *Weed Science* 56: 807–813.
  - Shivrain V.K., N.R. Burgos, D.R. Gealy, M.A. Sales, and Kenneth L Smith. 2009. Gene flow from weedy red rice (*Oryza sativa* L.) to cultivated rice and fitness of hybrids. *Pest Management Science* 65:1124–1129.
  - Shivrain, V.K., N.R. Burgos, R.C. Scott, E.E. Gbur, Jr., L.E. Estorninos, Jr., M.R. McClelland. 2010. Diversity of weedy red rice (*Oryza sativa* L.) in Arkansas, U.S.A. in relation to weed management. *Crop Protection* 29:721–730.
  - Singh, V., S. Singh, H. Black, V. Boyett, S. Basu, D. Gealy, E. Gbur, A. Pereira, R.C. Scott, A. Caicedo, N.R. Burgos. 2017. Introgression of Clearfield rice crop traits into weedy red rice outcrosses. *Field Crops Research* 207:13–23.
  - Smith, R. J., Jr. 1988. Weed thresholds in southern U.S. rice *Oryza sativa*. *Weed Technology* 2:232–241.
  - Suh, H.S., Y.I. Sato, H. Morishima. 1997. Genetic characterization of weedy rice (*Oryza sativa* L.) based on morpho-physiology, isozymes and RAPD markers. *Theoretical and Applied Genetics* 94:316-321.
  - Sweeney, M.T., M.J. Thomson, B.E. Pfeil, and S. McCouch. 2006. Caught red-handed: Rc encodes a basic helix-loop-helix protein conditioning red pericarp in rice. *The Plant Cell* 18:283–294.
  - Tan, L., X. Li, F. Liu, X. Sun, C. Li, Z. Zhu, Y. Fu, H. Cai, X. Wang, D. Xie, and C. Sun. 2008. Control of a key transition from prostrate to erect growth in rice domestication. *Nature Genetics* doi:10.1038/ng.197.
  - Vaughan, L.K., B.V. Ottis, A.M. Prazak-Havey, C.A. Bormans, C. Sneller, J.M. Chandler, and W.D. Park. 2001. Is all red rice found in commercial rice really *Oryza sativa*? *Weed Science* 49(4):468-476.

- Vidotto, F. and A. Ferrero. 2009. Interactions between weedy rice and cultivated rice in Italy. *Italian Journal of Agronomy*: 127-136.
- Vidotto, F. and A. Ferrero. 2000. Germination behaviour of red rice (*Oryza sativa L.*) seeds in field and laboratory conditions. *Agronomie, EDP Sciences* 20:375-382.
- Vigueira, C.C., Qi, X., Song, B.-K., Li, L.-F., Caicedo, A.L. et al., 2019. Call of the wild rice: *Oryza rufipogon* shapes weedy rice evolution in Southeast Asia. *Evolutionary Applications* 12:93–104.
- Wolfe, D.W., A.T. DeGaetano, G.M. Peck, M. Carey, L.H. Ziska, J. Lea-Cox, A.R. Kemanian, M.P. Hoffmann, and D.Y. Hollinger. 2018. Unique challenges and opportunities for northeastern US crop production in a changing climate. *Climatic Change* 146:231–245.
- Wu, Y., S. Zhao, X. Li, B. Zhang, L. Jiang, Y. Tang, J. Zhao, X. Ma, H. Cai, C. Sun, and L. Tan. 2018. Deletions linked to *PROG1* gene participate in plant architecture domestication in Asian and African rice. *Nature Communications* 9:4157.
- Young, M.L., J.K. Norsworthy, R.C. Scott, and L.T. Barber. 2017. Tolerance of Southern US Rice Cultivars to Benzobicyclon. *Weed Technology* 31:658-665.
- Young, M.L. 2017. Evaluation of benzobicyclon for use in Midsouthern rice (*Oryza sativa*) systems. MSc Thesis, University of Arkansas. 122 p.
- Zhu, B.-F., L. Si, Z. Wang, Y. Zhou, J. Zhu et al., 2011. Genetic control of a transition from black to straw-white seed hull in rice domestication. *Plant Physiology* 155:1301–1311.
- Ziska, L.H., D.R. Gealy, N. Burgos, A.L. Caicedo, J. Gressel, A.L. Lawton-Rauh, L.A. Avila, G. Theisen, J. Norsworthy, A. Ferrero, F. Vidotto, D.E. Johnson, F.G. Ferreira, E. Marchesan, V. Menezes, M.A. Cohn, S. Linscombe, L. Carmona, R. Tang, A. Merotto Junior. 2015. Weedy (red) rice: an emerging constraint to global rice production. *Advances in Agronomy* 129:181–228.
- Ziska, L.H., M.B. Tomecek, and D.R. Gealy. 2010b. Competitive interactions between cultivated and red rice as a function of recent and projected increases in atmospheric carbon dioxide. *Agronomy Journal* 102:118-123.

#### Ελληνόγλωσσον

- Ελευθεροχωρινός, Η.Γ. 2014. Αντιμετώπιση ζιζανίων στην καλλιέργεια του ρυζιού. *Γεωργία-Κτηνοτροφία* 10:112-121.
- Ελευθεροχωρινός, Η.Γ. 2017. Ρύζι: Επιλογή κατάλληλου ζιζανιοκτόνου. *Γεωργία-Κτηνοτροφία* 4:72-75.
- Ελευθεροχωρινός, Η.Γ. 2018. Η εφαρμογής της ψευδοσποράς για την αντιμετώπιση ζιζανίων σε ορυζώνες. *Γεωργία-Κτηνοτροφία* 3:66-67.
- Ελευθεροχωρινός, Η.Γ. 2019. Προβλήματα διαχείρισης ζιζανίων στην καλλιέργεια του ρυζιού: Παρατηρήσεις και συμπεράσματα από την περιοχή Θεσσαλονίκης κατά το 2019. *Γεωργία-Κτηνοτροφία* 9:70-73.